

ROMANIAN MATHEMATICAL MAGAZINE


In any $\triangle ABC$ with p_a, p_b, p_c

\rightarrow Spieker cevians, the following relationship holds :

$$\frac{\sqrt{p_a n_a}}{h_a} + \frac{\sqrt{p_b n_b}}{h_b} + \frac{\sqrt{p_c n_c}}{h_c} \geq \frac{\sqrt{6}}{3} \left(\frac{b+c}{a} + \frac{c+a}{b} + \frac{a+b}{c} \right) + 3 - 2\sqrt{6}$$

Proposed by Mohamed Amine Ben Ajiba-Tanger-Morocco

Solution by Soumava Chakraborty-Kolkata-India



Let AS produced meet BC at X and $m(\angle BAX) = \alpha$ and $m(\angle CAX) = \beta$ (say)
and inradius of $\triangle DEF = r'$ (say)

$$\text{Now, } 16[\triangle DEF]^2 = 2 \sum \left(\frac{a^2}{4} \right) \left(\frac{b^2}{4} \right) - \sum \frac{a^4}{16} = \frac{1}{16} \left(2 \sum a^2 b^2 - \sum a^4 \right) = \frac{16r^2 s^2}{16}$$

$$\Rightarrow [\triangle DEF] = \frac{rs}{4} \Rightarrow r' \left(\frac{\frac{a}{2} + \frac{b}{2} + \frac{c}{2}}{2} \right) = \frac{rs}{4} \Rightarrow r' = \frac{r}{2} \rightarrow (1)$$

$$\because \text{Spieker center is incenter of } \triangle DEF, \therefore m(\angle AFS) = B + \frac{C}{2} = \frac{2B + C}{2} = \frac{B + \pi - A}{2}$$

$$= \frac{\pi}{2} - \frac{A - B}{2} \text{ and } m(\angle AES) = C + \frac{B}{2} = \frac{\pi}{2} - \frac{A - C}{2} \rightarrow (2)$$

Via (1), (2) and using cosine law on $\triangle AFS$ and $\triangle AES$, we arrive at :

$$\begin{aligned} AS^2 &= \frac{r^2}{4\sin^2 \frac{C}{2}} + \frac{c^2}{4} - \left(\frac{2r}{2\sin \frac{C}{2}} \right) \left(\frac{c}{2} \right) \sin \frac{A-B}{2} \\ &= \frac{r^2}{4\sin^2 \frac{B}{2}} + \frac{b^2}{4} - \left(\frac{2r}{2\sin \frac{B}{2}} \right) \left(\frac{b}{2} \right) \sin \frac{A-C}{2} \\ \Rightarrow 2AS^2 &\stackrel{(1)}{=} \frac{r^2}{4\sin^2 \frac{C}{2}} + \frac{c^2}{4} - \left(\frac{2r}{2\sin \frac{C}{2}} \right) \left(\frac{c}{2} \right) \sin \frac{A-B}{2} + \frac{r^2}{4\sin^2 \frac{B}{2}} + \frac{b^2}{4} \\ &\quad - \left(\frac{2r}{2\sin \frac{B}{2}} \right) \left(\frac{b}{2} \right) \sin \frac{A-C}{2} \end{aligned}$$

ROMANIAN MATHEMATICAL MAGAZINE

Now, $\left(\frac{2r}{2\sin\frac{c}{2}}\right)\left(\frac{c}{2}\right)\sin\frac{A-B}{2} + \left(\frac{2r}{2\sin\frac{B}{2}}\right)\left(\frac{b}{2}\right)\sin\frac{A-C}{2}$

$$= \frac{r}{2} \left(4R\cos\frac{C}{2} \sin\frac{A-B}{2} + 4R\cos\frac{B}{2} \sin\frac{A-C}{2} \right)$$

$$= Rr \left(2\sin\frac{A+B}{2} \sin\frac{A-B}{2} + 2\sin\frac{A+C}{2} \sin\frac{A-C}{2} \right)$$

$$= Rr \left(1 - 2\sin^2\frac{B}{2} + 1 - 2\sin^2\frac{C}{2} - 2 \left(1 - 2\sin^2\frac{A}{2} \right) \right)$$

$$= 2Rr \left(\frac{2a(s-b)(s-c) - b(s-c)(s-a) - c(s-a)(s-b)}{abc} \right)$$

$$= \frac{Rr}{8Rrs} (2a^3 + (b+c)a^2 - 2a(b^2 + c^2) - (b+c)(b-c)^2)$$

$$= \frac{4(b+c)bcc\sin^2\frac{A}{2} - 2a \cdot 2bcc\cos A}{8s} = \frac{bc \left((2s-a)\sin^2\frac{A}{2} - a \left(1 - 2\sin^2\frac{A}{2} \right) \right)}{2s}$$

$$= \frac{bc \left((2s+a)\sin^2\frac{A}{2} - a \right)}{2s} = \frac{(2s+a)(s-b)(s-c)}{2s} - 2Rr$$

$$\Rightarrow - \left(\frac{2r}{2\sin\frac{c}{2}} \right) \left(\frac{c}{2} \right) \sin\frac{A-B}{2} - \left(\frac{2r}{2\sin\frac{B}{2}} \right) \left(\frac{b}{2} \right) \sin\frac{A-C}{2}$$

$$\stackrel{(*)}{=} \frac{-(2s+a)(s-b)(s-c)}{2s} + 2Rr$$

Again, $\frac{r^2}{4\sin^2\frac{B}{2}} + \frac{r^2}{4\sin^2\frac{C}{2}} = \frac{r^2}{4} \left(\frac{ca}{(s-c)(s-a)} + \frac{ab}{(s-a)(s-b)} \right)$

$$= \frac{r^2}{4r^2s} (ca(s-b) + ab(s-c)) = \frac{ab+ca}{4} - 2Rr \stackrel{(**)}{=} \frac{r^2}{4\sin^2\frac{B}{2}} + \frac{r^2}{4\sin^2\frac{C}{2}}$$

$$(i), (*), (**) \Rightarrow 2AS^2 = \frac{b^2 + c^2 + ab + ca}{4} - \frac{(2s+a)(s-b)(s-c)}{2s}$$

$$= \frac{(a+b+c)(b^2 + c^2 + ab + ca) - (2a+b+c)(c+a-b)(a+b-c)}{8s}$$

$$= \frac{b^3 + c^3 - abc + a(2b^2 + 2c^2 - a^2)}{4s} \Rightarrow 2AS^2 \stackrel{(ii)}{=} \frac{b^3 + c^3 - abc + a(4m_a^2)}{4s}$$

Via sine law on ΔAFS , $\frac{r}{2\sin\frac{c}{2}\sin\alpha} = \frac{AS}{\cos\frac{A-B}{2}} = \frac{4s}{(a+b)\sin\frac{c}{2}}$

$$\Rightarrow csin\alpha \stackrel{(***)}{=} \frac{r(a+b)}{2AS} \text{ and via sine law on } \Delta AES, bsin\beta \stackrel{((***)}{=} \frac{r(a+c)}{2AS}$$

Now, $[BAX] + [BAX] = [ABC] \Rightarrow \frac{1}{2}p_a csin\alpha + \frac{1}{2}p_a bsin\beta = rs$

via (***)) and (((**))) $\Rightarrow \frac{p_a(a+b+a+c)}{4AS} = s \Rightarrow p_a = \frac{4s}{2s+a} AS$

$$\Rightarrow p_a^2 \stackrel{\text{via (ii)}}{=} \frac{16s^2}{(2s+a)^2} \cdot \frac{b^3 + c^3 - abc + a(4m_a^2)}{8s}$$

ROMANIAN MATHEMATICAL MAGAZINE

$$\therefore p_a^2 \stackrel{(*)}{=} \frac{2s}{(2s+a)^2} (b^3 + c^3 - abc + a(4m_a^2))$$

$$\begin{aligned}
& \text{Now, } b^3 + c^3 - abc + a(4m_a^2) = b^3 + c^3 - abc + a(2b^2 + 2c^2 - a^2) \\
&= (b+c)(b^2 - bc + c^2) + a(b^2 - bc + c^2) + a(b^2 + c^2 - a^2) \\
&= 2s(b^2 - bc + c^2) + a(b^2 - bc + c^2 + bc - a^2) \\
&= (2s+a)(b^2 - bc + c^2) + a\left(\frac{(b+c)^2 - (b-c)^2}{4} - a^2\right) \\
&= (2s+a)(b^2 - bc + c^2) + \frac{a(b+c+2a)(b+c-2a)}{4} - \frac{a(b-c)^2}{4} \\
&= (2s+a)(b^2 - bc + c^2) + \frac{a(2s-a+2a)(b+c-2a)}{4} - \frac{a(b-c)^2}{4} \\
&= (2s+a) \cdot \frac{4b^2 + 4c^2 - 4bc + a(b+c-2a)}{4} - \frac{a(b-c)^2}{4} \\
&= (2s+a).
\end{aligned}$$

$$\frac{4(z+x)^2 + 4(x+y)^2 - 4(z+x)(x+y) + (y+z)((z+x)+(x+y)-2(y+z))}{4}$$

$$\begin{aligned}
& - \frac{a(b-c)^2}{4} \quad (a = y+z, b = z+x, c = x+y) \\
&= (2s+a) \cdot \frac{4x(x+y+z) + 2x(y+z) + 3(y-z)^2}{4} - \frac{a(b-c)^2}{4} \\
&= (2s+a) \left(s(s-a) + \frac{3}{4}(b-c)^2 + \frac{a(s-a)}{2} \right) - \frac{a(b-c)^2}{4} \\
&= (2s+a) \left(s(s-a) + \frac{3}{4}(b-c)^2 + \frac{a(s-a)}{2} \right) - \frac{(a+2s-2s)(b-c)^2}{4} \\
&= (2s+a) \left(s(s-a) + \frac{(b-c)^2}{2} + \frac{a(s-a)}{2} \right) + \frac{s(b-c)^2}{2}
\end{aligned}$$

$$\therefore b^3 + c^3 - abc + a(4m_a^2) \stackrel{(**)}{=} (2s+a) \left(\frac{(s-a)(2s+a)}{2} + \frac{(b-c)^2}{2} \right) + \frac{s(b-c)^2}{2}$$

$$\begin{aligned}
& \therefore (\bullet), (\bullet\bullet) \Rightarrow p_a^2 = \frac{2s}{(2s+a)^2} \left(\frac{(s-a)(2s+a)^2}{2} + \frac{(2s+a)(b-c)^2}{2} + \frac{s(b-c)^2}{2} \right) \\
&= s(s-a) + (b-c)^2 \left(\left(\frac{s}{2s+a} \right)^2 + \frac{s}{2s+a} + \frac{1}{4} - \frac{1}{4} \right) \\
&= s(s-a) - \frac{(b-c)^2}{4} + (b-c)^2 \cdot \left(\frac{s}{2s+a} + \frac{1}{2} \right)^2 \\
&= s(s-a) + \frac{(b-c)^2}{4} \left(\frac{(4s+a)^2}{(2s+a)^2} - 1 \right) \\
&\Rightarrow p_a^2 \stackrel{(***)}{=} s(s-a) + \frac{s(3s+a)(b-c)^2}{(2s+a)^2}
\end{aligned}$$

$$\text{Now, } p_a \geq \frac{2b^2 - bc + 2c^2}{6R} \Leftrightarrow \frac{p_a^2}{h_a^2} \geq \frac{(2b^2 - bc + 2c^2)^2}{36R^2 \cdot \frac{b^2c^2}{4R^2}}$$

ROMANIAN MATHEMATICAL MAGAZINE

$$\begin{aligned}
& \Leftrightarrow \frac{p_a^2 - h_a^2}{h_a^2} \geq \left(\frac{2b^2 - bc + 2c^2}{3bc} - 1 \right) \left(\frac{2b^2 - bc + 2c^2}{3bc} + 1 \right) \\
\text{via (***)} \quad & \Leftrightarrow \frac{s(s-a) + \frac{s(3s+a)(b-c)^2}{(2s+a)^2} - s(s-a) + \frac{s(s-a)(b-c)^2}{a^2}}{h_a^2} \geq \frac{4(b-c)^2(b^2 + bc + c^2)}{9b^2c^2} \\
& \Leftrightarrow \frac{4s^4(b-c)^2}{a^2h_a^2(2s+a)^2} \geq \frac{4(b-c)^2(b^2 + bc + c^2)}{9b^2c^2} \\
& \Leftrightarrow \frac{s^4}{4s(s-a)(s-b)(s-c)(2s+a)^2} \geq \frac{b^2 + bc + c^2}{9b^2c^2} \quad (\because (b-c)^2 \geq 0) \\
& \Leftrightarrow \frac{9s^3b^2c^2}{4(s-a)(2s+a)^2} \geq (b^2 + bc + c^2)(s-b)(s-c) \\
& \quad = (b^2 + bc + c^2)(-s(s-a) + bc) \\
& \Leftrightarrow \frac{9s^3b^2c^2}{4(s-a)(2s+a)^2} \geq (bc - s(s-a))(b^2 + c^2) + b^2c^2 - bcs(s-a) \\
& \Leftrightarrow \frac{9s^3b^2c^2}{4(s-a)(2s+a)^2} - b^2c^2 + bcs(s-a) \geq (bc - s(s-a))((2s-a)^2 - 2bc) \\
& \quad \Leftrightarrow \frac{9s^3b^2c^2}{4(s-a)(2s+a)^2} - b^2c^2 + bcs(s-a) \geq \\
& \quad (bc - s(s-a))(2s-a)^2 - 2b^2c^2 + 2bcs(s-a) \\
& \Leftrightarrow \frac{9s^3b^2c^2}{4(s-a)(2s+a)^2} + b^2c^2 + s(s-a)(2s-a)^2 - bc(s(s-a) + (2s-a)^2) \geq 0 \\
& \Leftrightarrow \boxed{\frac{25s^3 - 12sa^2 - 4a^3}{4(s-a)(2s+a)^2} \cdot b^2c^2 - (5s^2 - 5sa + a^2) \cdot bc + s(s-a)(2s-a)^2 \stackrel{(\square)}{\geq} 0}
\end{aligned}$$

Now, LHS of (□) is a quadratic polynomial with discriminant =

$$\begin{aligned}
& (5s^2 - 5sa + a^2)^2 - \frac{25s^3 - 12sa^2 - 4a^3}{(2s+a)^2} \cdot s(2s-a)^2 \\
& = \frac{-a^2(12s^4 - 18s^3a + 5s^2a^2 + 2sa^3 - a^4)}{(2s+a)^2} \\
& = \frac{-a^2(s-a)((s-a)(12s^2 + 6sa + 5a^2) + 6a^3)}{(2s+a)^2} < 0 \quad (\because s > a) \\
& \therefore (\square) \text{ is true (strict inequality)}
\end{aligned}$$

$$\therefore p_a \geq \frac{2b^2 - bc + 2c^2}{6R} \text{ and analogs } \rightarrow (m)$$

Again, Stewart's theorem $\Rightarrow b^2(s-c) + c^2(s-b) = an_a^2 + a(s-b)(s-c)$

$$\begin{aligned}
\Rightarrow s(b^2 + c^2) - bc(2s-a) &= an_a^2 + a(s^2 - s(2s-a) + bc) \Rightarrow s(b^2 + c^2) - 2sbc \\
&= an_a^2 + a(as - s^2) \Rightarrow s(b^2 + c^2 - a^2 - 2bc) = an_a^2 - as^2 \Rightarrow an_a^2 = as^2 + \\
&s(2bc \cos A - 2bc) = as^2 - 4sbc \sin^2 \frac{A}{2} = as^2 - \frac{4sbc(s-b)(s-c)(s-a)}{bc(s-a)} \\
&= as^2 - s(a^2 - (b-c)^2) = as(s-a) + s(b-c)^2 \\
&\Rightarrow n_a^2 = s(s-a) + \frac{s}{a}(b-c)^2 \rightarrow (l)
\end{aligned}$$

ROMANIAN MATHEMATICAL MAGAZINE

Now, $n_a \stackrel{?}{\geq} \frac{b^2 - bc + c^2}{2R} \Leftrightarrow \frac{n_a}{h_a} \stackrel{?}{\geq} \frac{b^2 - bc + c^2}{bc} \Leftrightarrow \frac{n_a^2}{h_a^2} - 1 \stackrel{?}{\geq} \left(\frac{b^2 - bc + c^2}{bc} \right)^2 - 1$

$$= \frac{(b-c)^2(b^2+c^2)}{b^2c^2} \Leftrightarrow \frac{n_a^2 - h_a^2}{h_a^2} \stackrel{?}{\geq} \frac{(b-c)^2(b^2+c^2)}{b^2c^2} \text{ via (l)} \Leftrightarrow$$

$$s(s-a) + \frac{s}{a}(b-c)^2 - s(s-a) + \frac{s(s-a)(b-c)^2}{a^2} \stackrel{?}{\geq} \frac{(b-c)^2(b^2+c^2)}{b^2c^2} \cdot \frac{b^2c^2}{4R^2}$$

$$\Leftrightarrow \left(\frac{s}{a} + \frac{s(s-a)}{a^2} \right) (b-c)^2 \stackrel{?}{\geq} \frac{(b-c)^2(b^2+c^2)}{4R^2} \Leftrightarrow \frac{s^2}{a^2} \stackrel{?}{\geq} \frac{b^2+c^2}{4R^2} (\because (b-c)^2 \geq 0)$$

$$\Leftrightarrow 4R^2s^2 \stackrel{?}{\geq} a^2b^2 + c^2a^2 \rightarrow \text{true (strict inequality)} \because 4R^2s^2 \stackrel{\text{Goldstone}}{\geq}$$

$$\sum_{\text{cyc}} a^2b^2 > a^2b^2 + c^2a^2 \therefore n_a \geq \frac{b^2 - bc + c^2}{2R} \text{ and analogs} \rightarrow (n)$$

$$\therefore \text{via (m) and (n), } \frac{p_a n_a}{h_a^2} \geq \frac{\frac{2b^2 - bc + 2c^2}{6R} \cdot \frac{b^2 - bc + c^2}{2R}}{\frac{b^2c^2}{4R^2}} \Rightarrow \frac{p_a n_a}{h_a^2} - 1 - \frac{2}{3} \cdot \frac{(b-c)^4}{b^2c^2}$$

$$\geq \frac{(2b^2 - bc + 2c^2)(b^2 - bc + c^2) - 3b^2c^2 - 2(b-c)^4}{3b^2c^2} = \frac{5bc(b-c)^2}{3b^2c^2}$$

$$\geq \frac{2\sqrt{6}(b-c)^2}{3bc} \therefore \frac{p_a n_a}{h_a^2} \geq 1 + \frac{2}{3} \cdot \frac{(b-c)^4}{b^2c^2} + \frac{2\sqrt{6}(b-c)^2}{3bc} = \left(1 + \frac{\sqrt{6}(b-c)^2}{3bc} \right)^2$$

$$\Rightarrow \frac{\sqrt{p_a n_a}}{h_a} \geq 1 + \frac{\sqrt{6}(b-c)^2}{3bc} \text{ and analogs} \therefore \frac{\sqrt{p_a n_a}}{h_a} + \frac{\sqrt{p_b n_b}}{h_b} + \frac{\sqrt{p_c n_c}}{h_c} \geq$$

$$3 + \frac{\sqrt{6}}{3} \cdot \sum_{\text{cyc}} \frac{b^2 + c^2 - 2bc}{bc} = 3 + \frac{\sqrt{6}}{3} \cdot \left(\sum_{\text{cyc}} \frac{b+c}{a} - 6 \right) \therefore \frac{\sqrt{p_a n_a}}{h_a} + \frac{\sqrt{p_b n_b}}{h_b} + \frac{\sqrt{p_c n_c}}{h_c} \geq$$

$$\geq \frac{\sqrt{6}}{3} \left(\frac{b+c}{a} + \frac{c+a}{b} + \frac{a+b}{c} \right) + 3 - 2\sqrt{6} \forall \Delta ABC,$$

with equality iff ΔABC is equilateral (QED)