

ROMANIAN MATHEMATICAL MAGAZINE


In any ΔABC with p_a, p_b, p_c

\rightarrow Spieker cevians, the following relationship holds :

$$\sqrt{p_a p_b} + \sqrt{p_b p_c} + \sqrt{p_c p_a} \geq \frac{2(a^2 + b^2 + c^2) + ab + bc + ca}{6R}$$

Proposed by Mohamed Amine Ben Ajiba-Tanger-Morocco

Solution by Soumava Chakraborty-Kolkata-India



Proof : Let AS produced meet BC at X and $m(\angle BAX) = \alpha$ and $m(\angle CAX) = \beta$ (say)
and inradius of $\Delta DEF = r'$ (say)

$$\begin{aligned} \text{Now, } 16[DEF]^2 &= 2 \sum \left(\frac{a^2}{4} \right) \left(\frac{b^2}{4} \right) - \sum \frac{a^4}{16} = \frac{1}{16} \left(2 \sum a^2 b^2 - \sum a^4 \right) = \frac{16r^2 s^2}{16} \\ \Rightarrow [DEF] &= \frac{rs}{4} \Rightarrow r' \left(\frac{\frac{a}{2} + \frac{b}{2} + \frac{c}{2}}{2} \right) = \frac{rs}{4} \Rightarrow r' = \frac{r}{2} \rightarrow (1) \end{aligned}$$

$$\begin{aligned} \because \text{Spieker center is incenter of } \Delta DEF, \therefore m(\angle AFS) &= B + \frac{C}{2} = \frac{2B + C}{2} = \frac{B + \pi - A}{2} \\ &= \frac{\pi}{2} - \frac{A - B}{2} \text{ and } m(\angle AES) = C + \frac{B}{2} = \frac{\pi}{2} - \frac{A - C}{2} \rightarrow (2) \end{aligned}$$

Via (1), (2) and using cosine law on ΔAFS and ΔAES , we arrive at :

$$AS^2 = \frac{r^2}{4\sin^2 \frac{C}{2}} + \frac{c^2}{4} - \left(\frac{2r}{2\sin \frac{C}{2}} \right) \left(\frac{c}{2} \right) \sin \frac{A-B}{2}$$

ROMANIAN MATHEMATICAL MAGAZINE

$$\begin{aligned}
 &= \frac{r^2}{4\sin^2 \frac{B}{2}} + \frac{b^2}{4} - \left(\frac{2r}{2\sin \frac{B}{2}} \right) \left(\frac{b}{2} \right) \sin \frac{A-C}{2} \\
 \Rightarrow 2AS^2 &\stackrel{(i)}{=} \frac{r^2}{4\sin^2 \frac{C}{2}} + \frac{c^2}{4} - \left(\frac{2r}{2\sin \frac{C}{2}} \right) \left(\frac{c}{2} \right) \sin \frac{A-B}{2} + \frac{r^2}{4\sin^2 \frac{B}{2}} + \frac{b^2}{4} \\
 &\quad - \left(\frac{2r}{2\sin \frac{B}{2}} \right) \left(\frac{b}{2} \right) \sin \frac{A-C}{2}
 \end{aligned}$$

$$\begin{aligned}
 \text{Now, } & \left(\frac{2r}{2\sin \frac{C}{2}} \right) \left(\frac{c}{2} \right) \sin \frac{A-B}{2} + \left(\frac{2r}{2\sin \frac{B}{2}} \right) \left(\frac{b}{2} \right) \sin \frac{A-C}{2} \\
 &= \frac{r}{2} \left(4R \cos \frac{C}{2} \sin \frac{A-B}{2} + 4R \cos \frac{B}{2} \sin \frac{A-C}{2} \right) \\
 &= Rr \left(2 \sin \frac{A+B}{2} \sin \frac{A-B}{2} + 2 \sin \frac{A+C}{2} \sin \frac{A-C}{2} \right) \\
 &= Rr \left(1 - 2 \sin^2 \frac{B}{2} + 1 - 2 \sin^2 \frac{C}{2} - 2 \left(1 - 2 \sin^2 \frac{A}{2} \right) \right) \\
 &= 2Rr \left(\frac{2a(s-b)(s-c) - b(s-c)(s-a) - c(s-a)(s-b)}{abc} \right) \\
 &= \frac{Rr}{8Rrs} (2a^3 + (b+c)a^2 - 2a(b^2 + c^2) - (b+c)(b-c)^2)
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{4(b+c)b \cos^2 \frac{A}{2} - 2a \cdot 2bc \cos A}{8s} = \frac{bc \left((2s-a) \sin^2 \frac{A}{2} - a \left(1 - 2 \sin^2 \frac{A}{2} \right) \right)}{2s} \\
 &= \frac{bc \left((2s+a) \sin^2 \frac{A}{2} - a \right)}{2s} = \frac{(2s+a)(s-b)(s-c)}{2s} - 2Rr \\
 &\Rightarrow - \left(\frac{2r}{2\sin \frac{C}{2}} \right) \left(\frac{c}{2} \right) \sin \frac{A-B}{2} - \left(\frac{2r}{2\sin \frac{B}{2}} \right) \left(\frac{b}{2} \right) \sin \frac{A-C}{2} \\
 &\stackrel{(*)}{=} \frac{-(2s+a)(s-b)(s-c)}{2s} + 2Rr
 \end{aligned}$$

$$\begin{aligned}
 \text{Again, } & \frac{r^2}{4\sin^2 \frac{B}{2}} + \frac{r^2}{4\sin^2 \frac{C}{2}} = \frac{r^2}{4} \left(\frac{ca}{(s-c)(s-a)} + \frac{ab}{(s-a)(s-b)} \right) \\
 &= \frac{r^2}{4r^2 s} (ca(s-b) + ab(s-c)) = \frac{ab+ca}{4} - 2Rr \stackrel{(**)}{=} \frac{r^2}{4\sin^2 \frac{B}{2}} + \frac{r^2}{4\sin^2 \frac{C}{2}} \\
 &\stackrel{(i), (*), (**)}{\Rightarrow} 2AS^2 = \frac{b^2 + c^2 + ab + ca}{4} - \frac{(2s+a)(s-b)(s-c)}{2s} \\
 &= \frac{(a+b+c)(b^2 + c^2 + ab + ca) - (2a+b+c)(c+a-b)(a+b-c)}{8s} \\
 &= \frac{b^3 + c^3 - abc + a(2b^2 + 2c^2 - a^2)}{4s} \Rightarrow 2AS^2 \stackrel{(ii)}{=} \frac{b^3 + c^3 - abc + a(4m_a^2)}{4s}
 \end{aligned}$$

ROMANIAN MATHEMATICAL MAGAZINE

Via sine law on ΔAFS , $\frac{r}{2\sin\frac{c}{2}\sin\alpha} = \frac{AS}{\cos\frac{A-B}{2}} = \frac{cAS}{(a+b)\sin\frac{c}{2}}$
 $\Rightarrow c\sin\alpha \stackrel{(***)}{=} \frac{r(a+b)}{2AS}$ and via sine law on ΔAES , $b\sin\beta \stackrel{****)}{=} \frac{r(a+c)}{2AS}$

Now, $[BAX] + [BAX] = [ABC] \Rightarrow \frac{1}{2}p_a c\sin\alpha + \frac{1}{2}p_a b\sin\beta = rs$

$$\begin{aligned} &\stackrel{\text{via } (***) \text{ and } ****)}{=} \frac{p_a(a+b+a+c)}{4AS} = s \Rightarrow p_a = \frac{4s}{2s+a} AS \\ &\Rightarrow p_a^2 \stackrel{\text{via (ii)}}{=} \frac{16s^2}{(2s+a)^2} \cdot \frac{b^3 + c^3 - abc + a(4m_a^2)}{8s} \\ &\therefore \boxed{p_a^2 \stackrel{(*)}{=} \frac{2s}{(2s+a)^2} (b^3 + c^3 - abc + a(4m_a^2))} \end{aligned}$$

$$\begin{aligned} \text{Now, } b^3 + c^3 - abc + a(4m_a^2) &= b^3 + c^3 - abc + a(2b^2 + 2c^2 - a^2) \\ &= (b+c)(b^2 - bc + c^2) + a(b^2 - bc + c^2) + a(b^2 + c^2 - a^2) \\ &= 2s(b^2 - bc + c^2) + a(b^2 - bc + c^2 + bc - a^2) \\ &= (2s+a)(b^2 - bc + c^2) + a\left(\frac{(b+c)^2 - (b-c)^2}{4} - a^2\right) \\ &= (2s+a)(b^2 - bc + c^2) + \frac{a(b+c+2a)(b+c-2a)}{4} - \frac{a(b-c)^2}{4} \\ &= (2s+a)(b^2 - bc + c^2) + \frac{a(2s-a+2a)(b+c-2a)}{4} - \frac{a(b-c)^2}{4} \\ &= (2s+a). \frac{4b^2 + 4c^2 - 4bc + a(b+c-2a)}{4} - \frac{a(b-c)^2}{4} \\ &= (2s+a). \end{aligned}$$

$$\boxed{4(z+x)^2 + 4(x+y)^2 - 4(z+x)(x+y) + (y+z)((z+x) + (x+y) - 2(y+z))}$$

$$\begin{aligned} &- \frac{a(b-c)^2}{4} \quad (a = y+z, b = z+x, c = x+y) \\ &= (2s+a). \frac{4x(x+y+z) + 2x(y+z) + 3(y-z)^2}{4} - \frac{a(b-c)^2}{4} \\ &= (2s+a) \left(s(s-a) + \frac{3}{4}(b-c)^2 + \frac{a(s-a)}{2} \right) - \frac{a(b-c)^2}{4} \\ &= (2s+a) \left(s(s-a) + \frac{3}{4}(b-c)^2 + \frac{a(s-a)}{2} \right) - \frac{(a+2s-2s)(b-c)^2}{4} \\ &= (2s+a) \left(s(s-a) + \frac{(b-c)^2}{2} + \frac{a(s-a)}{2} \right) + \frac{s(b-c)^2}{2} \end{aligned}$$

$$\therefore \boxed{b^3 + c^3 - abc + a(4m_a^2) \stackrel{(*)}{=} (2s+a) \left(\frac{(s-a)(2s+a)}{2} + \frac{(b-c)^2}{2} \right) + \frac{s(b-c)^2}{2}}$$

$$\begin{aligned} &\therefore (*) \Rightarrow p_a^2 = \frac{2s}{(2s+a)^2} \left(\frac{(s-a)(2s+a)^2}{2} + \frac{(2s+a)(b-c)^2}{2} + \frac{s(b-c)^2}{2} \right) \\ &= s(s-a) + (b-c)^2 \left(\left(\frac{s}{2s+a} \right)^2 + \frac{s}{2s+a} + \frac{1}{4} - \frac{1}{4} \right) \end{aligned}$$

ROMANIAN MATHEMATICAL MAGAZINE

$$\begin{aligned}
&= s(s-a) - \frac{(b-c)^2}{4} + (b-c)^2 \cdot \left(\frac{s}{2s+a} + \frac{1}{2} \right)^2 \\
&= s(s-a) + \frac{(b-c)^2}{4} \left(\frac{(4s+a)^2}{(2s+a)^2} - 1 \right) \\
&\Rightarrow p_a^2 \stackrel{(\dots)}{=} s(s-a) + \frac{s(3s+a)(b-c)^2}{(2s+a)^2} \\
\text{Now, } p_a &\geq \frac{2b^2 - bc + 2c^2}{6R} \Leftrightarrow \frac{p_a^2}{h_a^2} \geq \frac{(2b^2 - bc + 2c^2)^2}{36R^2 \cdot \frac{b^2 c^2}{4R^2}} \\
&\Leftrightarrow \frac{p_a^2 - h_a^2}{h_a^2} \geq \left(\frac{2b^2 - bc + 2c^2}{3bc} - 1 \right) \left(\frac{2b^2 - bc + 2c^2}{3bc} + 1 \right) \\
\text{via } (\dots) \quad &\Leftrightarrow \frac{s(s-a) + \frac{s(3s+a)(b-c)^2}{(2s+a)^2} - s(s-a) + \frac{s(s-a)(b-c)^2}{a^2}}{h_a^2} \geq \frac{4(b-c)^2(b^2 + bc + c^2)}{9b^2c^2} \\
&\Leftrightarrow \frac{4s^4(b-c)^2}{a^2 h_a^2 (2s+a)^2} \geq \frac{4(b-c)^2(b^2 + bc + c^2)}{9b^2c^2} \\
&\Leftrightarrow \frac{s^4}{4s(s-a)(s-b)(s-c)(2s+a)^2} \geq \frac{b^2 + bc + c^2}{9b^2c^2} \quad (\because (b-c)^2 \geq 0) \\
&\Leftrightarrow \frac{9s^3b^2c^2}{4(s-a)(2s+a)^2} \geq (b^2 + bc + c^2)(s-b)(s-c) \\
&\quad = (b^2 + bc + c^2)(-s(s-a) + bc) \\
&\Leftrightarrow \frac{9s^3b^2c^2}{4(s-a)(2s+a)^2} \geq (bc - s(s-a))(b^2 + c^2) + b^2c^2 - bcs(s-a) \\
&\Leftrightarrow \frac{9s^3b^2c^2}{4(s-a)(2s+a)^2} - b^2c^2 + bcs(s-a) \geq (bc - s(s-a))((2s-a)^2 - 2bc) \\
&\Leftrightarrow \frac{9s^3b^2c^2}{4(s-a)(2s+a)^2} - b^2c^2 + bcs(s-a) \geq \\
&\quad (bc - s(s-a))(2s-a)^2 - 2b^2c^2 + 2bcs(s-a) \\
&\Leftrightarrow \frac{9s^3b^2c^2}{4(s-a)(2s+a)^2} + b^2c^2 + s(s-a)(2s-a)^2 - bc(s(s-a) + (2s-a)^2) \geq 0 \\
&\Leftrightarrow \boxed{\frac{25s^3 - 12sa^2 - 4a^3}{4(s-a)(2s+a)^2} \cdot b^2c^2 - (5s^2 - 5sa + a^2) \cdot bc + s(s-a)(2s-a)^2 \stackrel{(\square)}{\geq} 0}
\end{aligned}$$

Now, LHS of (\square) is a quadratic polynomial with discriminant =

$$\begin{aligned}
&(5s^2 - 5sa + a^2)^2 - \frac{25s^3 - 12sa^2 - 4a^3}{(2s+a)^2} \cdot s(2s-a)^2 \\
&= \frac{-a^2(12s^4 - 18s^3a + 5s^2a^2 + 2sa^3 - a^4)}{(2s+a)^2} \\
&= \frac{-a^2(s-a)((s-a)(12s^2 + 6sa + 5a^2) + 6a^3)}{(2s+a)^2} < 0 \quad (\because s > a) \\
&\therefore (\square) \text{ is true (strict inequality)} \\
&\therefore p_a \geq \frac{2b^2 - bc + 2c^2}{6R} \text{ and analogs}
\end{aligned}$$

ROMANIAN MATHEMATICAL MAGAZINE

$$\begin{aligned}
 \therefore p_b p_c &\geq \frac{(2c^2 - ca + 2a^2)(2a^2 - ab + 2b^2)}{36R^2} \stackrel{?}{\geq} \frac{(4a^2 + 4bc - ab - ac)^2}{144R^2} \\
 \Leftrightarrow 15a^2(b - c)^2 &\stackrel{?}{\geq} 0 \rightarrow \text{true} \therefore p_b p_c \geq \frac{(4a^2 + 4bc - ab - ac)^2}{144R^2} \\
 \Rightarrow \boxed{\sqrt{p_b p_c} \geq \frac{4a^2 + 4bc - ab - ac}{12R}} \quad \text{and analogs} \therefore \sqrt{p_a p_b} + \sqrt{p_b p_c} + \sqrt{p_c p_a} \geq \\
 &\frac{1}{12R} \cdot \left(4 \sum_{\text{cyc}} a^2 + 4 \sum_{\text{cyc}} bc - \sum_{\text{cyc}} ab - \sum_{\text{cyc}} ac \right) \\
 \therefore \sqrt{p_a p_b} + \sqrt{p_b p_c} + \sqrt{p_c p_a} &\geq \frac{2(a^2 + b^2 + c^2) + ab + bc + ca}{6R} \\
 \forall \Delta ABC, " = " \text{ iff } \Delta ABC \text{ is equilateral (QED)}
 \end{aligned}$$